



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

It is merely subjective opinion that patient-reported outcome measures are not objective tools

Citation for published version:

Hamilton, DF, Giesinger, JM & Giesinger, K 2017, 'It is merely subjective opinion that patient-reported outcome measures are not objective tools', *Bone & Joint Research*, vol. 6, no. 12, pp. 665-666.
<https://doi.org/10.1302/2046-3758.612.BJR-2017-0347>

Digital Object Identifier (DOI):

[10.1302/2046-3758.612.BJR-2017-0347](https://doi.org/10.1302/2046-3758.612.BJR-2017-0347)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Bone & Joint Research

Publisher Rights Statement:

Attributions licence (CC-BY-NC), which permits unrestricted use, distribution, and reproduction in any medium, but not for commercial gain, provided the original author and source are credited.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.





■ EDITORIAL

It is merely subjective opinion that patient-reported outcome measures are not objective tools

**D. F. Hamilton,
J. M. Giesinger,
K. Giesinger**

*The British Editorial
Society of Bone and
Joint Surgery, London,
United Kingdom*

The last decade has seen a paradigm shift in the measurement of clinical outcomes, with an increasing focus placed on the patients' perspective to compliment and augment the clinicians' report, imaging and laboratory results.

Patient-Reported Outcome Measures (PROMs) can be simply described as the patients' report of the status of their health. This can range from an informal comment as to their symptoms to the completion of a validated questionnaire, which through an algorithm, allows an objective quantification of the response. The distinction between 'an informal comment' and 'validated questionnaire' is critically important. A number of well-validated self-reported questionnaires are available to assess a patient's general and joint-specific health status. These validated PROMs are central to orthopaedic research, clinical practice, quality control and benchmarking.¹⁻⁹

Validated PROMs, as with other routinely applied clinical tests, are designed to provide robust and meaningful measurements. For outcome questionnaires, this is determined by the three main psychometric properties: objectivity, reliability and validity.¹⁰⁻¹² Objectivity implies that the result of a measurement should not be influenced by the person taking that measurement. For questionnaires objectivity is usually high, as these are completed independently by the patient him/herself. Reliability is generally evaluated by demonstrating test-retest reproducibility. Validity (content, construct and criterion), tests the degree to which a PROM actually measures what it proposes to measure. When considering longitudinal measurements, the responsiveness of a PROM is also important, i.e. the ability to detect changes over time.¹³

These generic properties are applicable, and central, to any measurement tool that doctors base their clinical decisions on.¹⁴

Many clinicians though are less trusting of 'subjective PROMs' than they are of 'objective measurements' such as angles determined on a plain radiograph, a fracture classification, or a physical performance test.¹⁵⁻²² They prefer what they perceive to be a 'hard' or 'real' measurement over some 'psychologically overlaid opinion of a patient who may be grumpy after a bad night'. This belief though is not in line with the evidence.

Hahn et al²³ have analysed the degree of error in commonly performed clinicians' measurements and compared it with the degree of error inherent to various validated PROMs. Interestingly, the PROMs compare favourably to the measures that clinicians trust in. For example, the test-retest correlations for heart rate ($r = 0.68$) or diastolic blood pressure ($r = 0.63$) are modest, whereas the physical functioning scale of the SF-36 shows very high agreement ($r = 0.93$). The literature as a whole reports a range of measurement characteristics for both well-established clinical tools and PROMs, examples of more and less robust measures can be found for each.

The 'distrust' of PROMs may be explained by the relative newness of these metrics. As with all evaluations, one must ensure suitable tool selection, application and interpretation. For constructs such as pain or satisfaction, the patient's perception is the only source of information and therefore PROMs should be considered the gold-standard evaluation. The picture is more complicated for constructs such as physical function, where this can be assessed through both PROMs and direct observation. Correlating the time taken to complete a specific test of physical function (e.g. the timed-get-up-and-go) with the results of a physical function questionnaire may not find particularly robust agreement. This should not be surprising as the questionnaire does not try to measure the time taken to complete a task,

■ D. F. Hamilton, PhD, MCSP,
Research Fellow, University of
Edinburgh, Orthopaedics, FU413
Chancellors Building, 49 Little
France Crescent, Edinburgh, EH16
4SB, UK.

■ J. M. Giesinger, PhD, Research
Fellow, Innsbruck Institute of
Patient Centered Outcome
Research, 6020 Innsbruck, Austria.

■ K. Giesinger, MD, MSc,
Department of Orthopaedic
Surgery and Traumatology,
Senior Consultant Knee Surgeon,
Kantonsspital St Gallen,
St Gallen, Switzerland.

Correspondence should be sent to
D. F. Hamilton;
email: dhamilt4@exseed.ed.ac.uk

doi: 10.1302/2046-3758.612.
BJR-2017-0347

Bone Joint Res 2017;6:665-666.

but instead asks about the patients' perception of the difficulty of completing the task. As such we would expect to see a similar direction of change in the respective scores when measuring the effect of an intervention, but to expect the same result misunderstands that PROMs capture a different aspect of outcome than a performance test does.²⁴ Similarly, it is quite reasonable for a patient to be satisfied with the outcome of surgery but report a low score on a validated PROM.^{25,26} This does not suggest that the PROM is 'wrong', but that the construct (theme of questions) evaluated by that PROM is not particularly associated with the criteria that determines the patient's report of satisfaction.

Hopefully time and familiarity will enhance the selection, application and interpretation of PROMs. The validated patient outcome questionnaire is not a 'subjective' opinion but an 'objective' evaluation that quantifies the patient's pain, function or severity of disease as perceived by the patient. Assuming the PROM has been well constructed, it provides a robust measurement and therefore should be recognised as an objective tool.

References

- Gagnier JJ. Patient reported outcomes in orthopaedics. *J Orthop Res* 2017;35:2098-2108.
- Malak TT, Broomfield JA, Palmer AJ, et al. Surrogate markers of long-term outcome in primary total hip arthroplasty: A systematic review. *Bone Joint Res* 2016;5:206-214.
- Kleinlugtenbelt YV, Nienhuis RW, Bhandari M, et al. Are validated outcome measures used in distal radial fractures truly valid? A critical assessment using the COnsensus-based Standards for the selection of health Measurement INstruments (COSMIN) checklist. *Bone Joint Res* 2016;5:153-161.
- Poitras S, Wood KS, Savard J, Dervin GF, Beaulé PE. Predicting early clinical function after hip or knee arthroplasty. *Bone Joint Res* 2015;4:145-151.
- Hamilton DF, Ghert M, Simpson AH. Interpreting regression models in clinical outcome studies. *Bone Joint Res* 2015;4:152-153.
- Parsons N, Griffin XL, Achten J, Costa ML. Outcome assessment after hip fracture: is EQ-5D the answer? *Bone Joint Res* 2014;3:69-75.
- Hamilton DF, Giesinger JM, Patton JT, et al. Making the Oxford Hip and Knee Scores meaningful at the patient level through normative scoring and registry data. *Bone Joint Res* 2015;4:137-144.
- Barlow T, Griffin D, Barlow D, Realpe A. Patients' decision making in total knee arthroplasty: a systematic review of qualitative research. *Bone Joint Res* 2015;4:163-169.
- Kingsbury SR, Dube B, Thomas CM, Conaghan PG, Stone MH. Is a questionnaire and radiograph-based follow-up model for patients with primary hip and knee arthroplasty a viable alternative to traditional regular outpatient follow-up clinic? *Bone Joint J* 2016;98-B:201-208.
- Terwee CB, Bot SDM, de Boer MR, et al. Quality criteria were proposed for measurement properties of health status questionnaires. *J Clin Epidemiol* 2007;60:34-42.
- Mokkink LB, Terwee CB, Patrick DL, et al. The COSMIN study reached international consensus on taxonomy, terminology, and definitions of measurement properties for health-related patient-reported outcomes. *J Clin Epidemiol* 2010;63:737-745.
- Giesinger K, Hamilton DF, Jost B, Holzner B, Giesinger JM. Comparative responsiveness of outcome measures for total knee arthroplasty. *Osteoarthritis Cartilage* 2014;22:184-189.
- Hamilton DF, Giesinger JM, MacDonald DJ, et al. Responsiveness and ceiling effects of the Forgotten Joint Score-12 following total hip arthroplasty. *Bone Joint Res* 2016;5:87-91.
- Langlois J, Hamadouche M. Current recommendations for assessing the reliability of a measurement tool: a survival guide for orthopaedic surgeons. *Bone Joint J* 2016;98-B:166-172.
- Konan S, Hossain F, Patel S, Haddad FS. Measuring function after hip and knee surgery: the evidence to support performance-based functional outcome tasks. *Bone Joint J* 2014;96-B:1431-1435.
- Luna IE, Kehlert H, Peterson B, et al. Early patient-reported outcomes versus objective function after total hip and knee arthroplasty: a prospective cohort study. *Bone Joint J* 2017;99-B:1167-1175.
- Harrell K, Clark R, Downes K, Virani N, Frankle M. Correlation of subjective and objective measures before and after shoulder arthroplasty. *Orthopedics* 2013;36:808-814.
- Tawonsawatruk T, Hamilton DF, Simpson AHRW. Validation of fracture callus scoring systems in animal models. *J Orthop Res* 2014;32:1117-1119.
- Chopra S, Moerenhout K, Crevoisier. Subjective versus objective assessment in early clinical outcome of modified Lapidus procedure for hallux valgus deformity. *Clin Biomech (Bristol, Avon)* 2016;32:187-193.
- Toogood PA, Abdel MP, Spear JA, et al. The monitoring of activity at home after total hip arthroplasty. *Bone Joint J* 2016;98-B:1450-1454.
- Hossain FS, Konan S, Patel S, Rodriguez-Merchan EC, Haddad FS. The assessment of outcome after total knee arthroplasty: are we there yet? *Bone Joint J* 2015;97-B:3-9.
- Ramkumar PN, Harris JD, Noble PC. Patient-reported outcome measures after total knee arthroplasty: a systematic review. *Bone Joint Res* 2015;4:120-127.
- Hahn EA, Cella D, Chassany O, et al. Precision of health-related quality-of-life data compared with other clinical measures. *Mayo Clin Proc* 2007;82:1244-1254.
- Hamilton DF, Gaston P, Simpson AH. Is patient reporting of physical function accurate following total knee replacement? *J Bone Joint Surg [Br]* 2012;94-B:1506-1510.
- Hamilton DF, Lane JV, Gaston P, et al. What determines patient satisfaction with surgery? A prospective cohort study of 4709 patients following total joint replacement. *BMJ Open* 2013;3:e002525.
- Rogers BA, Alolabi B, Carrothers AD, Kreder HJ, Jenkinson RJ. Can the pre-operative Western Ontario and McMaster score predict patient satisfaction following total hip arthroplasty? *Bone Joint J* 2015;97-B:150-153.

Funding Statement

None declared

Author Contribution

- D. F. Hamilton: Writing the paper.
- J. M. Giesinger: Writing the paper.
- K. Giesinger: Writing the paper.

Conflicts of Interest Statement

None declared

© 2017 Hamilton, Giesinger and Giesinger. This is an open-access article distributed under the terms of the Creative Commons Attributions licence (CC-BY-NC), which permits unrestricted use, distribution, and reproduction in any medium, but not for commercial gain, provided the original author and source are credited.